

Appendix B, Chapter 14

Green Sturgeon

TABLE OF CONTENTS

14.0	Green Sturgeon (<i>Acipenser medirostris</i>).....	14-1
14.1	Life History & Requirements	14-1
14.2	Population Identification	14-2
14.3	Status & Abundance Trends.....	14-2
14.4	Limiting Factors	14-3
14.5	References	14-5
14.6	Other References of Interest.....	14-5



14.0 Green Sturgeon (*Acipenser medirostris*)

Green sturgeon is an anadromous species that spawn in several West Coast rivers but spend most of their life in near-shore marine and estuarine waters from Mexico to southeast Alaska (Houston 1988; Moyle *et al.* 1995). Significant spawning populations of green sturgeon have been identified in the Sacramento, Klamath, and Rogue Rivers. While green sturgeon do not spawn in the Columbia Basin, significant populations of subadults and adults are present in the estuary during summer and early fall. Green sturgeon are occasionally observed as far upriver as Bonneville Dam. Reasons for concentrations in the Columbia River are unclear because no spawning occurs in that system and all of the green sturgeon stomachs examined to date have been empty. These fish may be seeking warmer summer river waters in the northern part of their range.

NOAA Fisheries (formerly known as NMFS) completed a status review for green sturgeon in 2003 and determined that listing under the ESA was not warranted at this time.

14.1 Life History & Requirements

Adult green sturgeon typically migrate into fresh water beginning in late February (Moyle *et al.* 1995). Spawning occurs in deep turbulent river mainstems. Klamath and Rogue River populations appear to spawn within 100 miles of the ocean, while the Sacramento spawning run may travel over 200 miles. Spawning occurs from March–July, with peak activity from April–June (Moyle *et al.* 1995). Confirmed spawning populations in North America are in the Rogue, Klamath, and Sacramento Rivers (Moyle *et al.* 1995).

Specific spawning habitat preferences are unclear, but eggs likely are broadcast over large cobble where they settle into the cracks (Moyle *et al.* 1995). The adhesiveness of green sturgeon eggs is poor compared to white sturgeon (Van Eenennaam *et al.* 2001), which may be explained by the reduced thickness of the outer layer of the chorion of green sturgeon eggs (approximately half the thickness of that in white sturgeon; Deng *et al.* 2001). Optimum flow and temperature requirements for spawning and incubation are unclear, but spawning success in most sturgeons is related to these factors (Dettlaff *et al.* 1993). Temperatures above 68°F (20°C) were lethal to embryos in laboratory experiments (Cech *et al.* 2000).

Green sturgeon larvae are distinguished from other sturgeon by the absence of a swim-up or post-hatching pelagic stage. They can be distinguished from white sturgeon by their size (longer and larger), light pigmentation, and size and shape of the yolk-sac (Deng *et al.* 2001). Larvae hatched in the laboratory are photonegative, exhibiting hiding behavior (Deng *et al.* 2001), and after the onset of exogenous feeding, green sturgeon larvae and juveniles appear to be

nocturnal (Cech *et al.* 2000). This development pattern and behavior may be an adaptation suited for avoiding downstream displacement.

Juvenile green sturgeon grow rapidly reaching 2 feet within 2–3 years (Nakamoto and Kisanuki 1995). Juveniles appear to spend up from 1–4 years in fresh and estuarine waters and disperse into salt water at lengths of 1-2.5 feet. Green sturgeon are benthic feeders on invertebrates including shrimp and amphipods, small fish, and possibly mollusks (Houston 1988).




Figure 14-1. Conceptual model of green sturgeon life cycle and limiting factors

14.2 Population Identification

Based on a review of the limited genetic data available for green sturgeon, NMFS identified two distinct population segments. A northern DPS includes Klamath and Rogue spawning populations. A southern DPS includes the Sacramento spawning population. Fish using the Columbia River apparently include individuals from both population segments.

An Asian form of green sturgeon (*A. mikadoi*) also has been described and is known to spawn only in the Tumlin River (Artyukhin and Andronov 1990; Birstein et al 1997 as cited in Birstein and Bemis 1997).

14.3 Status & Abundance Trends

Available time series data were limited, but were not consistent with a modern decline in green sturgeon abundance. Time series data on green sturgeon abundance and size composition are limited to fishery landing statistics; these do not provide a consistent index of green sturgeon abundance. Columbia River harvest per unit effort and size composition data suggest an increasing rather than decreasing trend in green sturgeon abundance. Current data indicate that:

- green sturgeon still spawn in most systems where they were historically present,
- significant numbers of spawners are present in several systems, and
- geographic range of spawning green sturgeon is currently stable or increasing.

The wide distribution of green sturgeon, large numbers seasonally observed in some areas, and projections based on demographic rates suggest that total green sturgeon numbers are at least in the tens of thousands.

14.4 Limiting Factors

Spawning populations of green sturgeon are potentially affected by runoff and water management in natal rivers. Physical factors affecting green sturgeon habitat use in the lower Columbia River and estuary are unclear.

Like all sturgeon, green sturgeon populations are highly susceptible to fishing mortality. Significant catches or concentrations of green sturgeon have been reported in widely-scattered areas throughout the broad distribution of this species. Directed green sturgeon fisheries do not occur anywhere in their range, but small incidental harvest occurs in several areas. Annual harvest data are available from the Columbia River where salmon and white sturgeon fisheries intercept green sturgeon that gather in the estuary in large numbers during late summer and early fall. The majority of the green sturgeon harvest occurs in commercial gill net fisheries. Green sturgeon landings in the Columbia River declined steadily from a peak harvest of 6,400 in 1986, which occurred during expanded salmon fisheries for a very large fall chinook and coho return (Figure 14-2). Rather than an index of green sturgeon abundance, decreases are the direct result of increasing restrictions on white sturgeon and salmon fisheries since 1986.

Like the white sturgeon, green sturgeon probably recovered slowly following collapse of excessive 19th century fisheries. Gradual recovery is consistent with harvest patterns of green sturgeon in Columbia River fisheries: green sturgeon landings were identified beginning in 1938 and they show a generally increasing trend until the 1960s with variable harvest but no obvious trend after 1960. Recent fluctuations in green sturgeon landings likely result from a combination of annual variation in occurrence in the Columbia estuary and in salmon or white sturgeon fisheries. Where total harvest is expressed as catch per unit effort based on fish tickets logged by commercial fish buyers, catch rates have been variable and possibly increasing over the same period that total green sturgeon landings have decreased. Increases might have been even more pronounced but for the fact that the legal-size slot for retention was reduced from 48-72 inches to 48-66 inches TL in 1993. In addition, average size and frequency of large green sturgeon have been increasing.

No supplementation or conservation hatchery programs currently exist for green sturgeon throughout their range (Joel VanEnennaam, University of California, Davis, personal communication).




Figure 14-2. Recent annual harvest of green sturgeon as reported by Farr et al. (2002). Columbia includes Oregon and Washington sport and commercial fishery harvests. Klamath includes Yurok and Hoopa subsistence fishery harvests. Total also includes Oregon coastal sport, Washington coastal sport, commercial, Tribal, and Oregon and Washington trawl. The total does not include California sport harvest which is believed to be minor (Schaffter 2000).

14.5 References

- Artyukhin, E.N. and A.E. Andronov. 1990. A morphobiological study of the green sturgeon, *Acipenser medirostris* (Chondrostei, Acipenseridae), from the Tummin (Datta) River and some aspects of the ecology and zoogeography of the Acipenseridae. *Journal of Ichthyology* 30:11-21.
- Birstein, V. J. and W. E. Bemis. 1997. How many species are within the genus *Acipenser*? *Environmental Biology of Fishes* 48:157-163.
- Birstein, V.J., R. Hanner, R. DeSalle. 1997. Phylogeny of the Acipenseriformes: cytogenetic and molecular approaches. *Env. Bio of Fishes* 48:127-155.
- Cech, J.J., Jr., S.I. Doroshov, G.P. Moberg, B.P. May, R.G. Schaffter, and D.M. Kohlhorst. 2000. Biological assessment of green sturgeon in the Sacramento-San Joaquin watershed (Phase 1). Final Report to CALFED Bay-Delta Program (Project #98-C-15, Contract #B-81738).
- Deng, X., J. P. Van Eenennaam*, and Serge I. Doroshov. 2001. Comparison of early life stages and growth of green and white sturgeon.
- Detlaff, T. A., A. S. Ginsburg, and O. J. Schmalhausen. 1993. Sturgeon fishes: developmental biology and aquaculture. Springer-Verlag, Berlin 300 pp.
- Farr, R.A., M.L. Hughes, and T.A. Rien. 2002. Green sturgeon population characteristics in Oregon. Oregon Department of Fish and Wildlife Fish Research Project Annual Report F-178-R.
- Houston, J. J. 1988. Status of the green sturgeon, *Acipenser medirostris*, in Canada. *The Canadian Field Naturalist* 102:286-290.
- Moyle, P. B., R. M. Yoshiyama, J. E. Williams, and E. D. Wikramanayake. 1995. Green sturgeon. Pages 26-34 in *Fish species of Special Concern in California*, 2nd edition. Final Report to the Department of Fish and Game (contract 2128IF).
- Nakamoto, R.J., and T.T. Kisanuki. 1995. Age and growth of Klamath River green sturgeon (*Acipenser medirostris*). U.S. Fish and Wildlife Service Report 93-FP-13. Yreka, Ca.
- Schaffter, R.G. 2000. California Department of Fish and Game unpublished data in Beamesderfer (2000).
- Van Eenennaam, J. P., and S. I. Doroshov. 2001. Reproductive conditions of Klamath River green sturgeon (*Acipenser medirostris*). *Proceedings of International Symposium on Sturgeon, Wisconsin*. (In review for *Journal of Applied Ichthyology*).
- Van Eenennaam, J.P., M.A.H. Webb, X. Deng, S.I. Doroshov, R.B. Mayfield, J.J. Cech, Jr., D.C. Hillemeier, and T.E. Wilson. 2001. Artificial spawning and larval rearing of Klamath River green sturgeon. *Transactions of the American Fisheries Society* 130:159-165.

14.6 Other References of Interest

- Anders, P. J. 2000. Ancient fish need modern protection. *Fisheries* 25 (9):30.
- Artyukhin, E. N. 1998. The nature and origin of summer spawning sturgeons. *Voprosy Ikhtologii* 5:717-723 (in Russian, English translation in *Journal of Ichthyology* 5:717-723).

- Beamesderfer, R. C. P. and R. A. Farr. 1997. Alternatives for the protection and restoration of sturgeons and their habitat. *Environmental Biology of Fishes* 48:407-417.
- Beamesderfer, R. C. P., T. A. Rien, and A. A. Nigro. 1995. Differences in the dynamics and potential production of impounded and unimpounded white sturgeon populations in the lower Columbia River. *Transactions of the American Fisheries Society* 124:857-872.
- Bemis, W. E., and B. Kynard. 1977. Sturgeon rivers: an introduction to Acipenseriform biogeography and life history. *Environmental Biology of Fishes* 48:167-183.
- Bemis, W.E., V. J. Birstein, and J. R. Waldman. 1997a. Sturgeon biodiversity and conservation: an introduction. Pages 13-14, In: V. J. Birstein, J. R. Waldman, and W. E. Bemis eds., *Sturgeon Biodiversity and Conservation*. Kluwer Academic Publishers Dordrecht, Netherlands.
- Bemis, W. E., E. K. Findeis, and L. Grand. 1997b. An overview of Acipenseriformes. *Environmental Biology of Fishes* 48: 25-71.
- Bennett, D.H., T.J. Dresser, T.S. Curet, K.B. Lepla, and M.A. Madsen. 1993a. Lower Granite Reservoir in-water disposal test: Results of the fishery, benthic and habitat monitoring program Year-3 (1990). US Army Corps of Engineers, Walla Walla, Washington.
- Bennett, D.H., T.J. Dresser, T.S. Curet, K.B. Lepla, and M.A. Madsen. 1993b. Lower Granite Reservoir in-water disposal test: Results of the fishery, benthic and habitat monitoring program-Year 4 (1991). Completion Report. US Army Corps of Engineers, Walla Walla, Washington.
- Birstein, V.J. 1993. Sturgeons and paddlefishes: threatened fishes in need of conservation. *Conservation Biology* 7:773-787.
- Birstein, V.J. 1993. Is *Acipenser medirostris* one or two species? *The Sturgeon Quarterly* 1(2):8.
- Birstein, V. J., W. E. Bemis, and J. R. Waldman. 1997b. The threatened status of acipenseriform fishes: a summary. 1997. *Environmental Biology of Fishes* 48:427-435.
- Birstein, V. J., J. R. Waldman, and W. E. Bemis. 1997. *Sturgeon biodiversity and conservation*. Kluwer Academic Publishers. Dordrecht, The Netherlands.
- Brown, J. R. 1991. Molecular evolution and population genetics of sturgeon (Genus *Acipenser*) based on mitochondrial analysis. Ph.D. Thesis, Simon Fraser University, Burnaby, British Columbia.
- Bradford, M. J. 1992. Precision of recruitment predictions from early life stages of marine fishes. *Fishery Bulletin* 90:439-453.
- Brannon E., M. Powell, T. Quinn, and A. Talbot. 2002. Population structure of Columbia River Basin Chinook salmon and steelhead trout. Center for Salmonid and Freshwater Species at Risk, University of Idaho. Final report to the Bonneville Power Administration, Portland, Oregon. 178 pp.
- Brannon, E. L., K. P. Currens, D. Goodman, J. A. Lichatowich, B.E. Riddell, and R. N. Williams. W. E. MConaha, chair. 1999. Review of artificial anadromous and resident fish production in the Columbia River basin. Part I: A scientific basis for Columbia River production programs. NWPPC Document 99-4. Scientific Review team, Independent Scientific Advisory Board, Northwest Power Planning Council, Portland, Oregon. 132 pp.

- Brown, J. R., K. Beckenbach, A. T. Beckenbach, and M. J. Smith. 1996. Length variation, heteroplasmy and sequence divergence in the mitochondrial DNA of four species of sturgeon (Acipenser). *Genetics* 142:525-535.
- Bruch, R., F. P. Binkowski, and S. I. Doroshov, eds. 2001. Proceedings of the Fourth International Symposium on Sturgeons, Oshkosh, Wisconsin, USA, 8-13 July 2001.
- Buell, J. W. 1992. Fish entrainment monitoring of the Western-Pacific dredge RW Lofgren during operations outside the preferred work period. Buell and Associates, Inc. Portland, Oregon.
- Cooke, D. W., S. D. Leach, and J. J. Isely. Behavior and lack of upstream passage of shortnose sturgeon at a hydroelectric facility/navigation lock complex. Pages 101-110 In: W. VanWinkle, P. Anders, D. Dixon, and D. Secor, eds. *Biology, Management and Protection of North American Sturgeons*. American Fisheries Society Press, Bethesda, Maryland, USA.
- Craig, J. A. and R. L. Hacker. 1940. Sturgeon fisheries of the Columbia River Basin. *Bulletin of the Bureau of Fisheries* 49:204:208.
- de Vlaming, V. L. 1972. Environmental control of teleost reproductive cycles: a brief review. *Journal of Fish Biology* 4:131-140.
- Findeis, E. K. 1997. Osteology and phylogenetic interrelationships of sturgeons (Acipenseridae). *Environmental Biology of Fishes* 48:73-126.
- Grande, L., and W. E. Bemis. 1991. Osteology and phylogenetic relationships of fossil and recent paddlefishes (Polyodontidae) with comments on the interrelationships of Acipenseriformes. *Journal of Vertebrate Paleontology*, II supplement 1:1-121.
- Grande, L., and W. E. Bemis. 1997. A comprehensive phylogenetic study of amiid fishes (Amiidae) based on comparative skeletal anatomy. *Journal of Vertebrate Paleontology*.
- Ha, D. S. 2002. E-mail to R. Sitts, Metropolitan Water District of Southern California, Sacramento, California.
- Hart, J. L. 1973. Pacific Fishes of Canada. *Bulletin of the Fisheries Research Board of Canada* 180:83-84.
- Hillemeier, D. 2000. Yurok Tribe unpublished data in Beamesderfer (2000).
- Hjort, J. 1926. Fluctuations in the year class of important food fishes. *Journal du Conseil* 1:5-39.
- Houde, E. D. 1987. Fish early life dynamics and recruitment variability. Pages 17-29 in R. D. Hoyt, editor. 10th Annual Larval Fish Conference. American Fisheries Society Symposium 2. Bethesda, Maryland.
- Jin, F. 1995. Late Mesozoic acipenseriformes (Osteichthyes: Actinopterygii) in Central Asia and their biogeographical implications. Pages 15-21 In: A. Sun and Y. Yang, eds., *Sixth Symposium on Mesozoic Terrestrial Ecosystems and Biota, Short papers*, China Ocean Press, Beijing, China.
- Kempinger, J. J. 1988. Spawning and early life history of lake sturgeon in the Lake Winnebago System. *American Fisheries Society Symposium* 5:110-122.
- Klyashtorin, L. B. 1976. The sensitivity of young sturgeons to oxygen deficiency. *Journal of Ichthyology* 16:677-682.

- Kynard, B. 1997. Life history, latitudinal patterns, and status of shortnose sturgeon, *Acipenser brevirostrum*. *Environmental Biology of Fishes* 48:319-334.
- Lichtowich, J. A. 1999. *Salmon without rivers*. Island Press, Washington, DC. 317 pp.
- Long, J. A. 1995. *The rise of fishes*. Johns Hopkins University Press, Baltimore, Maryland.
- Lord, J. K. 1866. *The naturalist in Vancouver Island and British Columbia*. Volume 1(358p) and Volume 2 (375p). London.
- May, R. C. 1974. Larval mortality in marine fishes and the critical period concept. In J. H. S. Blaxter, editor. *The early life history of fish*. Springer-Verlag, New York.
- McDonald, M. 1894. Notes on the fisheries of the Pacific Coast, sturgeon and the sturgeon fishery.
- Migdalski, E. C. 1962. *Anglers guide to the freshwater sport fishes*. Ronald Press, New York.
- Munro, A. D., A. P. Scott, and T. J. Lam. 1990. *Reproductive seasonality in teleosts: environmental influences*. CRC Press, Inc. Boca Raton, Florida.
- Moyle, P. B. 1976. *Inland Fishes of California*. University of California Press, Berkeley, California. 405 pp.
- Moyle, P. B., and J. J. Cech, Jr. 1982. *Fishes: An Introduction to Ichthyology*. Prentice-Hall Publishers, Inc. Englewood Cliffs, NJ., USA.
- Musick, J. A., and 17 coauthors. 2000. Marine, estuarine, and diadromous fish stocks at risk of extinction in North America (exclusive of pacific salmonids). *Fisheries* 25(11)6-30.
- National Research Council (NRC). 1996. *Upstream: Salmon and society in the Pacific Northwest*. National Academy Press, Washington, DC.
- Oregon Department of Fish and Wildlife (ODFW). 1996. K. Beiningen and T. Rien, eds. *Effects of mitigative measures on productivity of white sturgeon populations in the Columbia River downstream from McNary Dam, and Determine status and habitat requirements of white sturgeon from the Columbia and Snake rivers upstream from McNary Dam*. Annual Progress Report to the Bonneville Power Administration. Project 86-50, Portland, Oregon.
- Rice, J. A., L. B. Crowder, and F. P. Binkowski. 1987. Evaluating potential sources of mortality for larval bloater: starvation and vulnerability to predation. *Canadian Journal of Fisheries and Aquatic Sciences* 44:467-472.
- Rochard, E. G., G. Castelnau, and M. Lepage. 1990. Sturgeons (Pisces: Acipenseridae): threats and prospects. *Journal of Fish Biology* 37 (Supplement A): 123-132.
- Scott, W.B. and E. J. Crossman. 1973. *Freshwater Fishes of Canada*. Bulletin 184. Fisheries Research Board of Canada.
- Secor, D., H. and Gunderson. 1988. Effects of hypoxia and temperature on survival, growth, and respiration of juvenile Atlantic sturgeon *Acipenser oxyrinchus*. *Fisheries Bulletin* 96:603-613.
- Secor, D. H., and E. J. Niklitschek. 2001. *Hypoxia and sturgeons: Report to the Chesapeake Bay Program dissolved oxygen criteria tema*. Technical report Series No. TS-314-01-CBL; Chesapeake Biological Laboratory, Solomons, MD.

- Secor, D. H., P. J. Anders, W. Van Winkle, and D. A. Dixon. 2002. Can We Study Sturgeons to Extinction? What We Do and Don't Know about the Conservation of North American Sturgeons. Pages 3-12 In: W. VanWinkle, P. Anders, D. Dixon, and D. Secor, eds. Biology, Management and Protection of North American Sturgeons. American Fisheries Society Symposium 28.
- Simpson, J., and R. Wallace. 1982. Fishes of Idaho. University of Idaho Press, Moscow, Idaho.
- Sinclair, M. 1988. Marine Populations: an essay on population regulation and speciation. University of Washington Press, Seattle and London. 252 pp.
- Stevens, D. E., and L. W. Miller. 1970. Distribution of sturgeon larvae in the Sacramento-San Joaquin river system. California Fish and Game 83:1-20.
- Tuell, M. A., and S. R. Everett. 2001. Evaluation of potential means of rebuilding sturgeon populations in the Snake River between Lower Granite and Hells Canyon dams. 2000 Annual Report, Preliminary Draft. Report to Bonneville Power Administration, Portland, Oregon. Contract No. 97-AM-30423, Project No. 00000333-00023 (9700900).
- Van Winkle, W., P. J. Anders, D. H. Secor, and D. A. Dixon, eds. 2002. Biology, Management, and Protection of Sturgeons. American Fisheries Society Symposium 28. 258 pp.
- Waldman, J. R. 1995. Sturgeon and paddlefishes: A convergence of biology, politics, and greed. Fisheries 20(9):20-49.
- Wang, Y. L., F. P. Binkowski, and S. I Doroshov. 1985. Effect of temperature on early development of white and lake sturgeon *Acipenser transmontanus*, and *A. fulvescens*. Environmental Biology of Fishes 14:43-50.
- Wright, S. 1931. Evolution in Mendelian populations. Genetics 16:97-159.
- Wydowski, R. S., and R. R. Whitney. 1979. Inland Fishes of Washington. University of Washington Press. Seattle.